

# Measure Theory Third Week

**Theorem (1.4.10):**

Let  $A$  be a Lebesgue measurable subset of  $\mathbf{R}$  such that  $\lambda(A) > 0$ .

The set  $\text{diff}(A) := \{x - y \mid x, y \in A\}$  contains an open interval containing 0.

**Proof:** By inner regularity, we can assume that  $A$  is compact.

With  $\lambda(A) = r > 0$ ,

there is an open set  $B$  such that  $B$  contains  $A$  and  $\lambda(B) < (1 + \epsilon)r$ . for any  $\epsilon > 0$ .

We require that  $\epsilon$  be less than 1.

As  $C := \mathbf{R} \setminus B$  is closed, disjoint from  $A$

and thus has a positive distance  $d$  to the compact set  $A$ ,

$A + \delta$  is contained in  $B$  for all  $\delta$  satisfying  $|\delta| < d$ .

But if there were no overlap between the sets  $A$  and  $A + \delta$  for any  $|\delta| < d$ ,

then by translation invariance  $A \cup (A + \delta)$  would be a Lebesgue measurable set of measure  $2r$  inside of  $B$ ,

which is impossible since  $\lambda(B) < (1 + \epsilon)r$ .

So for any given  $\delta$  with  $|\delta| < d$  there is an  $a \in A \cap A + \delta$ ,

meaning that  $a = a' + \delta$  for some other  $a' \in A$  and  $\delta = a - a'$ .  $\square$

We see that for every  $\epsilon$  there is a  $d$  such that all but an  $\epsilon$  fraction of the set  $A$  is used to get the difference set to include  $(-d, d)$ .

**Theorem (1.4.11):** Assuming the Axiom of Choice, there is a partition of  $\mathbf{R}$  into two parts  $A, B$ ,

meaning  $A \cap B = \emptyset$  and  $A \cup B = \mathbf{R}$ ,

such that for every finite interval  $I$  :

$$\lambda^*(A \cap I) = \lambda^*(B \cap I) = \lambda^*(I) \text{ and}$$

every Lebesgue measurable subset  $C$  either contained in either  $A$  or  $B$  has measure zero.

**Note:** The natural idea, a ring homomorphism from  $\mathbf{R}$  to  $\mathbf{Z}_2$  and letting  $A = \phi^{-1}(0)$  and  $B = \phi^{-1}(1)$ , is not possible,

whenever  $\phi(r) = 1$  then what should be  $\phi(\frac{r}{2})$ ?

Need a group homomorphism.

## Proof:

Let  $W = \mathbf{Q} + \mathbf{Z}\sqrt{2}$ ,

$\phi : W \rightarrow \mathbf{Z}_2$  is defined by

$$\phi\left(\frac{a}{b} + n\sqrt{2}\right) = n \pmod{2}.$$

Because  $\sqrt{2}$  is irrational,  $\phi$  is well defined and a group homomorphism:

$$\frac{a}{b} + n\sqrt{2} = \frac{a'}{b'} + n'\sqrt{2} \Rightarrow \sqrt{2} = \frac{a'}{b'(n-n')} - \frac{a}{b(n-n')} \in \mathbf{Q}, \text{ a contradiction when } n \neq n'.$$

Also both  $G_0 := \phi^{-1}(0) \subset W$  and  $G_1 := \phi^{-1}(1) \subset W$  are dense in  $\mathbf{R}$ , (and this can be shown with the Euclidean algorithm on the pair 1 and  $\sqrt{2}$  via smaller and smaller ways to write  $\frac{a}{b} + n\sqrt{2}$  with both even and odd  $n$ ).

Define an equivalence relation  $\sim$  by

$r \sim s$  if and only if  $r - s \in W$ .

Let  $E$  be a set such that  $|E \cap C| = 1$  (via Axiom of Choice) for every equivalence class  $C$ .

For every  $r \in \mathbf{R}$ ,  $r = e + \frac{a}{b} + n\sqrt{2}$ ,

for some  $e \in E$ ,  $a, b \in \mathbf{Z}$ ,  $n \in \mathbf{Z}$ , and uniquely so.

$A$  is the subset where  $n$  used is even and  $B$  is the subset where  $n$  used is odd.

$A$  and  $B$  are well defined because  $r$  cannot equal  $e' + \frac{a'}{b'} + n'\sqrt{2}$  for any other choices,

as then  $e - e'$  would be in  $W$  and  $e$  and  $e'$  would belong to the same equivalence class.

Assume that either  $A$  or  $B$  contained a Lebesgue measurable set of positive measure.

Either  $A - A$  or  $B - B$  must contain an open interval and hence some member of the dense set  $G_1$ , in other words

$$\frac{a_0}{b_0} + n_0\sqrt{2} = e_1 + \frac{a_1}{b_1} + n_1\sqrt{2} - e_2 - \frac{a_2}{b_2} - n_2\sqrt{2}$$

with  $n_0$  odd, both  $n_1$  and  $n_2$  either even or odd, and  $e_1, e_2 \in E$ .

As  $e_1$  and  $e_2$  must be equal (otherwise they would represent the same equivalence relation),  $n_0 = n_1 - n_2$  would be a contradiction.



Now suppose that either  $A \cap I$  or  $B \cap I$  has an outer Lebesgue measure less than  $I$  for some finite interval  $I$ .

That means  $A \cap I$  or  $B \cap I$  can be covered by some open set of measure strictly less than  $I$ .

implying that either  $I \setminus A = I \cap B$  or  $I \setminus B = I \cap A$  contains a closed set of positive measure, which, by the above, neither does.  $\square$

The same is true for three or more sets, but is much more difficult to show.

A measure  $\mu$  of a measure space  $(X, \mathcal{A}, \mu)$  is *complete*

if  $A \in \mathcal{A}$ ,  $\mu(A) = 0$  and  $B \subseteq A$  imply that  $B \in \mathcal{A}$ .

With  $(X, \mathcal{A}, \mu)$  a measure space,

the completion  $\mathcal{A}_\mu$  is the collection of subsets  $A$

for which there are sets  $E, F \in \mathcal{A}$

with  $E \subseteq A \subseteq F$  and  $\mu(F \setminus E) = 0$ .

The completion  $\bar{\mu}$  is the measure defined on  $\mathcal{A}_\mu$

such that  $\bar{\mu}(A) = \mu(E) = \mu(F)$ .

This is well defined as there cannot be two such levels (otherwise monotonicity is violated):

Suppose  $\mu(F') = \mu(E') > \mu(E) = \mu(F)$  with  $E' \subseteq A \subseteq F'$ ,  $E \subseteq A \subseteq F$ ,  $\mu(F' \setminus E') = \mu(F \setminus E) = 0$  with all four sets in  $\mathcal{A}$ .

It follows that  $E' \subseteq A \subseteq F$  and by containment  $\mu(E') \leq \mu(F)$ , a contradiction.

**Lemma (1.5.1):** Let  $(X, \mathcal{A}, \mu)$  be a measure space.

$\mathcal{A}_\mu$  is a  $\sigma$ -algebra on  $X$  that includes  $\mathcal{A}$

and  $\bar{\mu}$  is a measure defined on  $\mathcal{A}_\mu$  that is complete.

**Proof:** Containment of  $\mathcal{A}$  in  $\mathcal{A}_\mu$  and closure by complementation are trivial.

If  $A_1, A_2, \dots$  is a sequence of sets in  $\mathcal{A}_\mu$

and  $E_i$  and  $F_i$  are sequences in  $\mathcal{A}$

with  $\forall i \ E_i \subseteq A_i \subseteq F_i$  and  $\mu(F_i \setminus E_i) = 0$

then by countable additivity

$$0 = \sum_{i=1}^{\infty} \mu(F_i \setminus E_i) \geq \mu(\cup_{i=1}^{\infty} (F_i \setminus E_i)) \geq \mu(\cup_{i=1}^{\infty} F_i \setminus \cup_{i=1}^{\infty} E_i) \geq 0,$$

implying that  $\cup_{i=1}^{\infty} A_i \in \mathcal{A}_\mu$ .

And if the  $A_1, A_2, \dots$  are disjoint the same pairs  $E_i$  and  $F_i$  of sequences show that

$$\sum_{i=1}^{\infty} \mu(F_i) = \sum_{i=1}^{\infty} \mu(E_i) \leq \bar{\mu}(\cup_{i=1}^{\infty} A_i) \leq \sum_{i=1}^{\infty} \mu(F_i),$$

hence equality and countable additivity.  $\square$

Let  $(X, \mathcal{A}, \mu)$  be a measure space,  
and  $A$  any subset of  $X$ .

$$\mu^*(A) = \inf\{\mu(B) \mid A \subseteq B, B \in \mathcal{A}\} \text{ and}$$

$$\mu_*(A) = \sup\{\mu(B) \mid A \supseteq B, B \in \mathcal{A}\} .$$

$\mu^*(A)$  is the outer measure and  $\mu_*(A)$  is the inner measure.

**Lemma:**  $\mu^*$  is an outer measure.

**Proof:**  $\mu^*(\emptyset) = 0$  and monotonicity are trivial.

Let  $A_1, A_2, \dots$  be a sequence of sets.

Suppose that  $\sum_{i=1}^{\infty} \mu^*(A_i) < \infty$ :

For every  $i = 1, 2, \dots$  let  $B_i$  be a set in  $\mathcal{A}$  containing  $A_i$

such that  $\mu(B_i) \leq \mu^*(A_i) + \frac{\epsilon}{2^i}$ .

$B = \cup_{i=1}^{\infty} B_i$  includes  $A = \cup_{i=1}^{\infty} A_i$  and

$$\sum_{i=1}^{\infty} \mu^*(A_i) \geq \sum_{i=1}^{\infty} \mu(B_i) - \epsilon \geq \mu(B) - \epsilon \geq \mu^*(A) - \epsilon.$$

True for every  $\epsilon$  implies the inequality.  $\square$

**Lemma (1.5.5)** Given that  $\mu^*(A) < \infty$ ,  $A$  belongs to  $\mathcal{A}_\mu$  if and only if  $\mu_*(A) = \mu^*(A)$ .

**Proof:**  $\Rightarrow$  If  $A$  belongs to  $\mathcal{A}_\mu$  then there are sets  $E, F \in \mathcal{A}$  such that  $E \subseteq A \subseteq F$  and  $\mu(F \setminus E) = 0$ .

From  $\mu(E) \leq \mu_*(A) \leq \mu^*(A) \leq \mu(F)$

all are equal.



$\Leftarrow$  On the other hand, if  $\mu_*(E) = \mu^*(E) < \infty$

there are sequences of sets  $A_1, A_2, \dots$  and  $B_1, B_2, \dots$

with  $A_i \subseteq E$  and  $E \subseteq B_i$  and

$$\mu(A_i) \geq \mu_*(E) - \frac{1}{2^i} \text{ and } \mu^*(E) + \frac{1}{2^i} \geq \mu(B_i).$$

The sets  $A = \cup_i^\infty A_i$  and  $B = \cap_i^\infty B_i$

are both in  $\mathcal{A}$  and have the same common measure size  $\mu^*(E) = \mu_*(E)$ , with  $A \subseteq E \subseteq B$ .